



<https://helda.helsinki.fi>

Age-specific incidence of allergic and non-allergic asthma

Pakkasela, Johanna

2020-01-10

Pakkasela , J , Ilmarinen , P , Honkamäki , J , Tuomisto , L E , Andersen , H , Piirilä , P , Hisinger-Mölkänen , H , Sovijärvi , A , Backman , H , Lundbäck , B , Rönmark , E , Kankaanranta , H & Lehtimäki , L 2020 , ' Age-specific incidence of allergic and non-allergic asthma ' , BMC Pulmonary Medicine , vol. 20 , no. 1 , 9 . <https://doi.org/10.1186/s12890-019-1040-2>

<http://hdl.handle.net/10138/313356>

<https://doi.org/10.1186/s12890-019-1040-2>

cc_by

publishedVersion

Downloaded from Helda, University of Helsinki institutional repository.

This is an electronic reprint of the original article.

This reprint may differ from the original in pagination and typographic detail.


Please cite the original version.

RESEARCH ARTICLE

Open Access



Age-specific incidence of allergic and non-allergic asthma

Johanna Pakkasela^{1,2*} , Pinja Ilmarinen³, Jasmin Honkamäki², Leena E. Tuomisto³, Heidi Andersén⁴, Päivi Piirilä⁵, Hanna Hisinger-Mölkänen⁶, Anssi Sovijärvi⁵, Helena Backman⁷, Bo Lundbäck⁸, Eva Rönmark⁷, Hannu Kankaanranta^{2,3†} and Lauri Lehtimäki^{2,9†}

Abstract

Background: Onset of allergic asthma has a strong association with childhood but only a few studies have analyzed incidence of asthma from childhood to late adulthood in relation to allergy. The purpose of the study was to assess age-specific incidence of allergic and non-allergic asthma.

Methods: Questionnaires were sent to 8000 randomly selected recipients aged 20–69 years in Finland in 2016. The response rate was 52.3% ($n = 4173$). The questionnaire included questions on e.g. atopic status, asthma and age at asthma diagnosis. Asthma was classified allergic if also a physician-diagnosed allergic rhinitis was reported.

Results: The prevalence of physician-diagnosed asthma and allergic rhinitis were 11.2 and 17.8%, respectively. Of the 445 responders with physician-diagnosed asthma, 52% were classified as allergic and 48% as non-allergic. Median ages at diagnosis of allergic and non-allergic asthma were 19 and 35 years, respectively. Among subjects with asthma diagnosis at ages 0–9, 10–19, 20–29, 30–39, 40–49, 50–59 and 60–69 years, 70, 62, 58, 53, 38, 19 and 33%, respectively, were allergic. For non-allergic asthma, the incidence rate was lowest in children and young adults (0.7/1000/year). It increased after middle age and was highest in older age groups (2.4/1000/year in 50–59 years old).

Conclusions: The incidence of allergic asthma is highest in early childhood and steadily decreases with advancing age, while the incidence of non-allergic asthma is low until it peaks in late adulthood. After approximately 40 years of age, most of the new cases of asthma are non-allergic.

Keywords: Asthma, Allergic, Non-allergic, Adult, Child, Adolescent, Incidence, Prevalence, Onset, Age-specific

Background

Rackemann was the first to introduce the concept of extrinsic/allergic and intrinsic/non-allergic asthma in 1947 and thus described the first phenotypes of asthma [1]. Over recent decades, cluster analyses have confirmed that asthma is more of a heterogeneous disorder rather than just a single disease. Several phenotypes have been introduced in addition to the ones established 70 years ago [2–6], but differentiating the phenotypes in clinical practice can be challenging. One of the answers appears to be the age of asthma onset and subsequently the

division into childhood/early-onset and adult/late-onset asthma [6, 7].

Childhood asthma is commonly associated with allergy [8, 9]. Large cohorts have shown allergic sensitization as a risk factor for development [10, 11] and persistence of asthma in childhood [12]. On the other hand, there is a lack of comprehensive studies on the relevance of allergy to adult asthma. Although allergic sensitization has been reported as a risk factor for asthma in adults [13, 14] and adult-onset asthma [15], adult asthma is more often non-allergic than allergic [7, 16]. Also, the rates of allergic sensitization in adult-onset asthma are mostly below 50% [15–17]. According to a recent Finnish cluster analysis, allergic asthma diagnosed in adulthood was often associated with respiratory symptoms already during childhood [5]. However, contradicting results do also exist and a U.S.-based study reported only a slight

* Correspondence: johanna.pakkasela@tuni.fi

[†]Hannu Kankaanranta and Lauri Lehtimäki contributed equally to this work.

¹Department of Respiratory Medicine, Tampere University Hospital, FIN-33521 Tampere, Finland

²Faculty of Medicine and Health Technology, Tampere University, Tampere, Finland

Full list of author information is available at the end of the article



difference in allergic sensitization in early- and late-onset asthma (72 and 63% in subjects with asthma onset before and after 40 years of age, respectively) [18].

It appears that there is a lack of knowledge on the allergic and non-allergic phenotypes of adult-onset asthma and the relation between allergy and asthma onset age. Our aim was to study the association between asthma onset age and allergy by assessing age at diagnosis and age-specific incidence of asthma in adult asthmatics with and without allergic rhinitis in a population-based postal questionnaire study conducted in Finland.

Methods

Study design and population

The present study is part of the FinEsS (Finland-Estonia-Sweden) study, which is a postal questionnaire study on respiratory epidemiology conducted in collaboration in these three Northern European countries. Similar postal surveys were conducted in 1996, 2006 and 2016. The present study sample is part of the latest survey conducted in Finland in February 2016 and is formed from a random sample of 8000 subjects aged 20–69 years from the population in western Finland (Hospital Districts of Vaasa and Seinäjoki). The study sample was obtained from the Finnish Population Register and it was matched to the age and gender distribution of the population in the geographical area of our study. Finland is a bilingual country and the registered native language of a subject determined whether questionnaire in Finnish or Swedish language was used. The questionnaire was sent to a random sample of 7986 subjects after exclusion of subjects with unknown address. Two reminders were sent to those not responding. The sample size was 7942 subjects after further exclusion of subjects with non-analyzable data as shown in Fig. 1. In total, 4173 subjects responded yielding to a response rate of 52.3%. Of the responders, 206 were excluded because of missing data regarding smoking habits and thus, the actual sample size was 3967 responders included in the study. The study protocol was approved by the Ethical Committee of Helsinki University Hospital (approval number 200/13/03/00/15).

Study area

The study area is mainly rural with two major towns (Seinäjoki and Vaasa, about 62,000 and 68,000 inhabitants, respectively). It has a subarctic climate and the yearly average temperature is 4 °C (from −7 °C in the winter to 17 °C in the summer) [19]. The most common allergic sensitizations in Finland are against dogs, cats and pollens whereas sensitization to house dust mites and molds is less common [20].

Questionnaire and definitions

The FinEsS questionnaire was developed from the Obstructive Lung Disease in Northern Sweden (OLIN) questionnaire, which is modified from the Swedish translation of the British Medical Research Council (BMRC) questionnaire [21]. The questionnaire includes questions on respiratory diseases, symptoms, medication and comorbidities, risk factors and occupational factors considered relevant in respiratory epidemiology.


A physician-diagnosed asthma was defined by an answer “yes” to the question “Have you been diagnosed by a doctor as having asthma?”. Age at asthma diagnosis was determined by an answer to the question “What age were you when asthma was diagnosed?”. Allergic rhinitis was defined by an answer “yes” to either of the questions “Have you been diagnosed by a doctor as having allergic rhinitis caused by pollens (caused by e.g. birch, grass, mugwort)?” or “Have you been diagnosed by a doctor as having other allergic rhinitis (caused by e.g. cat or dog, but not pollen)?”. Allergic conjunctivitis was defined by an answer “yes” to the question “Have you been diagnosed by a doctor as having symptoms of allergy in your eyes?”. Age at diagnosis of allergy was not asked for. We used the presence and absence of allergic rhinitis as an indication of asthma being allergic or non-allergic, respectively. A sensitivity analysis was made by using the presence of either allergic rhinitis, allergic conjunctivitis or both as an indication of allergic asthma. Current smokers were considered those who reported smoking currently or during the 12 months preceding the survey. Ex-smokers reported previous smoking but had quit smoking at least 1 year prior to the survey. Never smokers did not report current or previous smoking.

Statistical analysis

Reconstructing age-specific incidence of asthma from cross-sectional data

Incidence of asthma in different age groups was estimated based on cross-sectional data on current age of the responders and age at diagnosis of asthma [22, 23]. Longitudinal data were retrospectively reconstructed from the questionnaire data as if the 3967 subjects were a cohort of newborns recruited between 69 and 20 years ago. A “time-to-event” (age at diagnosis of asthma) was recorded for each individual, and the population at risk at each age was updated by subtracting both events (subjects reporting asthma diagnosed at younger age) and censorships (asthma-naïve responders younger than the age for which population at risk was calculated) from the total sample.

In brief, subjects were divided into 10-year age groups based on their current age, and annual incidence of asthma per 1000 person-years (new asthma diagnoses/1000/year) was calculated in each age group by dividing



the number of incident asthmas in each group by age-group-specific population at risk, dividing the result by 10 and further multiplying with 1000. The 10-year age group -specific population at risk was a mean value of annually calculated respective 10-year risks. With respect to age 0, population at risk was all responders. For ages 1–20 years, subjects reporting asthma diagnosed at younger age than the age for which the population at

risk was calculated were subtracted to form the 1-year population at risk. The youngest responders were 20 years of age at the time of the study. After age 20, the responders who did not report physician-diagnosed asthma (i.e. asthma-naïve responders) and who were younger than the age for which the population at risk was calculated, were also subtracted from all responders to calculate populations at risk for ages 21–69 years.

7. Miranda C, Busacker A, Balzar S, Trudeau J, Wenzel SE. Distinguishing severe asthma phenotypes: role of age at onset and eosinophilic inflammation. *J Allergy Clin Immunol*. 2004;113:101–8.
8. Sears MR, Burrows B, Flannery EM, Herbison GP, Hewitt CJ, Holdaway MD. Relation between airway responsiveness and serum IgE in children with asthma and in apparently normal children. *N Engl J Med*. 1991;325:1067–71.
9. Sears MR, Burrows B, Flannery EM, Herbison GP, Holdaway MD. Atopy in childhood. I. Gender and allergen related risks for development of hay fever and asthma. *Clin Exp Allergy*. 1993;23:941–8.
10. Ulrik CS, Backer V, Hesse B, Dirksen A. Risk factors for development of asthma in children and adolescents: findings from a longitudinal population study. *Respir Med*. 1996;90:623–30.
11. Arshad SH, Tariq SM, Matthews S, Hakim E. Sensitization to common allergens and its association with allergic disorders at age 4 years: a whole population birth cohort study. *Pediatrics*. 2001;108:e33.
12. Sears MR, Greene JM, Willan AR, Wiecek EM, Taylor DR, Flannery EM, et al. A longitudinal, population-based, cohort study of childhood asthma followed to adulthood. *N Engl J Med*. 2003;349:1414–22.
13. Schoefer Y, Schäfer T, Meisinger C, Wichmann HE, Heinrich J. Predictivity of allergic sensitization (RAST) for the onset of allergic diseases in adults. *Allergy*. 2008;63:81–6.
14. Rönmark EP, Ekerljung L, Mincheva R, Sjölander S, Hagstad S, Wennergren G, et al. Different risk factor patterns for adult asthma, rhinitis and eczema: results from West Sweden asthma study. *Clin Transl Allergy*. 2016;6:28.
15. Rönmark E, Andersson C, Nyström L, Forsberg B, Järholm B, Lundbäck B. Obesity increases the risk of incident asthma among adults. *Eur Respir J*. 2005;25:282–8.
16. Amelink M, de Nijs SB, de Groot JC, van Tilburg PM, van Spiegel PI, Krouwels FH, et al. Three phenotypes of adult-onset asthma. *Allergy*. 2013;68:674–80.
17. Anto JM, Sunyer J, Basagana X, Garcia-Esteban R, Cerveri I, de Marco R, et al. Risk factors of new-onset asthma in adults: a population-based international cohort study. *Allergy*. 2010;65:1021–30.
18. Busse PJ, Cohn RD, Salo PM, Zeldin DC. Characteristics of allergic sensitization among asthmatic adults older than 55 years: results from the National Health and nutrition examination survey, 2005–2006. *Ann Allergy Asthma Immunol*. 2013;110:247–52.
19. Climatological statistics in Finland 1981–2010. The Finnish Meteorological Institute. <https://en.ilmatietaenlaitos.fi/statistics-from-1961-onwards>. Accessed 20 May 2018.
20. Pallasaho P, Rönmark E, Haahtela T, Sovijärvi AR, Lundbäck B. Degree and clinical relevance of sensitization to common allergens among adults: a population study in Helsinki, Finland. *Clin Exp Allergy*. 2006;36:503–9.
21. Lundbäck B, Nyström L, Rosenhall L, Stjernberg N. Obstructive lung disease in northern Sweden: respiratory symptoms assessed in a postal survey. *Eur Respir J*. 1991;4:257–66.
22. de Marco R, Locatelli F, Sunyer J, Burney P. Differences in incidence of reported asthma related to age in men and women. A retrospective analysis of the data of the European respiratory health survey. *Am J Respir Crit Care Med*. 2000;162:68–74.
23. Honkamäki J, Hisinger-Mölkänen H, Ilmarinen P, Piirilä P, Tuomisto LE, Andersén H, et al. Age- and gender-specific incidence of new asthma diagnosis from childhood to late adulthood. *Respir Med*. 2019;154:56–62.
24. Sergeant ESG. EpiTools epidemiological calculators: Ausvet Pty Ltd.; 2017. <http://epitools.ausvet.com.au>
25. Roorda RJ. Prognostic factors for the outcome of childhood asthma in adolescence. *Thorax*. 1996;51(Suppl 1):7–12.
26. Martinez FD. Links between pediatric and adult asthma. *J Allergy Clin Immunol*. 2001;107(Suppl):449–55.
27. Sood A, Qualls C, Schuyler M, Arynchyn A, Alvarado JH, Smith LJ, et al. Adult-onset asthma becomes the dominant phenotype among women by age 40 years. The longitudinal CARDIA study. *Ann Am Thorac Soc*. 2013;10:188–97.
28. Kankaanranta H, Tuomisto LE, Ilmarinen P. Age-specific incidence of new asthma diagnoses in Finland. *J Allergy Clin Immunol Pract*. 2017;5:189–91.e3.
29. Warm K, Hedman L, Lindberg A, Lötvall J, Lundbäck B, Rönmark E. Allergic sensitization is age-dependently associated with rhinitis, but less so with asthma. *J Allergy Clin Immunol*. 2015;136:1559–65.e2.
30. Siroux V, Basagaña X, Boudier A, Pin I, Garcia-Aymerich J, Vesin A, et al. Identifying adult asthma phenotypes using a clustering approach. *Eur Respir J*. 2011;38:310–7.
31. Jamrozik E, Knuiman MW, James A, Divitini M, Musk AW. Risk factors for adult-onset asthma: a 14 year longitudinal study. *Respirology*. 2009;14:814–21.
32. Romanet-Manent S, Charpin D, Magnan A, Lanteaume A, Vervloet D. Allergic vs nonallergic asthma: what makes the difference? *Allergy*. 2002;57:607–13.
33. Szczeklik A, Stevenson DD. Aspirin-induced asthma: advances in pathogenesis, diagnosis, and management. *J Allergy Clin Immunol*. 2003;111:913–21.
34. Warm K, Backman H, Lindberg A, Lundbäck B, Rönmark E. Low incidence and high remission of allergic sensitization among adults. *J Allergy Clin Immunol*. 2012;129:136–42.
35. Backman H, Räisänen P, Hedman L, Stridsman C, Andersson M, Lindberg A, et al. Increased prevalence of allergic asthma from 1996 to 2006 and further to 2016 – results from three population surveys. *Clin Exp Allergy*. 2017;47:1426–35.
36. Pallasaho P, Juusela M, Lindqvist A, Sovijärvi A, Lundbäck B, Rönmark E. Allergic rhinoconjunctivitis doubles the risk for incident asthma – results from a population study in Helsinki, Finland. *Respir Med*. 2011;105:1449–56.
37. Johansson SG, Hourihane JO, Bousquet J, Bruijnzeel-Koomen C, Dreborg S, Haahtela T, et al. A revised nomenclature for allergy: an EAACI position statement from the EAACI nomenclature task force. *Allergy*. 2001;56:813–24.
38. Burgess JA, Walters EH, Byrnes GB, Matheson MC, Jenkins MA, Wharton CL, et al. Childhood allergic rhinitis predicts asthma incidence and persistence to middle age: a longitudinal study. *J Allergy Clin Immunol*. 2007;120:863–9.
39. Guerra S, Sherrill DL, Martinez FD, Barbee RA. Rhinitis as an independent risk factor for adult-onset asthma. *J Allergy Clin Immunol*. 2002;109:419–25.
40. Global Initiative for Asthma. Global Strategy for Asthma Management and Prevention, 2018. <https://ginasthma.org/gina-reports/>. Accessed 15 May 2018.
41. Burgess JA, Walters EH, Byrnes GB, Wharton C, Jenkins MA, Abramson MJ, et al. Who remembers whether they had asthma as children? *J Asthma*. 2006;43:727–30.
42. Torén K, Palmqvist M, Löwhagen O, Balder B, Tunsäter A. Self-reported asthma was biased in relation to disease severity while reported year of asthma onset was accurate. *J Clin Epidemiol*. 2006;59:90–3.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
-